# Global warming - physicist's perspective - 04

Szymon P. Malinowski

University of Warsaw, Faculty of Physics, Institute of Geophysics





- Time-line (milestones) (after http://www.aip.org/history/climate/timeline.htm and other sources):
- Herschel hypothesizes on the effects of variablee solar emission om climate
- Fourier calculates that the Earth would be far colder if it lacked an atmosphere and introduces term "greenhouse effect"
- Tyndall discovers that some gases absorb infrared radiation. He suggests that changes in the concentration of the gases could bring climate change.
- Arrhenius publishes first calculation of global warming from increased concentrations of CO2.
- **1897** Chamberlin produces a model for global carbon exchange including feedbacks.
- 1930s: Global warming trend since late 19th century reported.

  Milankovitch proposes orbital changes as the cause of ice ages.

  Hulburt publishes calculations of warming from increased concentrations of CO2.
- Callendar argues that CO2 greenhouse global warming is underway, reviving interest in the problem.

The Stood Soriety of London

#### PHILOSOPHICAL TRANSACTIONS

hisas n

Part of Service of Service

.....

nia nea.

[ 265 ]

XIII. Observations tending to investigate the Nature of the Sun, in order to find the Causes or Symptoms of its variable Emission of Light and Heat; with Remarks on the Use that may possibly be drawn from Solar Observations. By William Herschel, L. L. D. F. R. S.

#### Read April 16, 1801.

On a former occasion I have shewn, that we have great reason to look upon the sun as a most magnificent habitable globe; and, from the observations which will be related in this Paper, it will now be seen, that all the arguments we have used before are not only confirmed, but that we are encouraged to go a considerable step farther, in the investigation of the physical and planetary construction of the sun. The influence of this eminent body, on the globe we inhabit, is so great, and so widely diffused, that it becomes almost a duty for us to study the operations which are carried on upon the solar surface. Since light and heat are so essential to our well-being, it must certainly be right for us to look into the source from whence they are derived, in order to see whether some material advantage may not be drawn from a thorough acquaintance with the causes from which they originate.

A similar motive engaged the Egyptians formerly to study and watch the motions of the Nile; and to construct instruments for measuring its rise with accuracy. They knew very well, that it was not in their power to add a single inch to the MDCCCI. M m



Observations Tending to Investigate the Nature of the Sun, in Order to Find the Causes or Symptoms of Its Variable Emission of Light and Heat; With Remarks on the Use That May Possibly Be Drawn from Solar Observations

#### William Herschel

Philosophical Transactions of the Royal Society of London Vol. 91 (1801), pp. 265-318



Fourier J (1827). "Mémoire Sur Les Températures Du Globe Terrestre Et Des Espaces Planétaires". Mémoires de l'Académie Royale des Sciences 7: 569–604

# **MÉMOIRES**

DE L'INSTITUT

DE FRANCE.

TOME VII.



PARIS,

CHEZ FIRMIN DIDOT, PÈRE ET FILS, LIBRAIRES,

RUE JACOB, 9° 24.

1827.

## MÉMOIRE

S CO BL

LES TEMPÉRATURES DU GLOBE TERRESTRE ET DES ESPACES PLANÉTAIRES.

PAR M. FOURIER

L'a question des températures terrestres. l'une des plus importantes et des plus difficiles de toute la philosophie naturelle, se compose d'éléments assez divers qui doivent être considérés sous un point de vue général. L'ai pense qu'il serait utile de réunir dans un seul écrit les conséquences principales de cette théorie; les détails analytiques que l'on omet-ici se trouvent pour la plupart dans les ouvrages que l'ai déja publiés. J'ai désiré surtout présentes aux physiciens, dans un tablesu peu étendu ; l'ensemble des phénomènes et les rapports mathématiques qu'ils ont entre eux mathématiques qu'ils entre eux m

La chaleur du globe terrestre dérive de trois sources qu'il est d'abord nécessaire de distinguée.

- 1º La terre est échauffée par les rayous solaires, dont l'inégale distribution produit la diversité des climats.
- 2º Elle participe à la température commune des espaces planétaires , étant exposée à l'irradiation des astres innombrables qui environment de toutes parts le système solaire.

Two basic climate mechanisms were known in the beginning of XIX century !!!



Eunice Newton Foote (July 17, 1819 – September 30, 1888)

was an American scientist, inventor, and women's rights campaigner. She was the first scientist to conclude that certain gases warmed when exposed to sunlight, and that rising carbon dioxide (CO2) levels would change atmospheric temperature and could affect climate, a phenomenon now referred to as the greenhouse effect.

#### ART. XXXI.—Circumstances affecting the Heat of the Sun's Rays; by EUNICE FOOTE.

#### (Read before the American Association, August 23d, 1856.)

My investigations have had for their object to determine the different circumstances that affect the thermal action of the rays of light that proceed from the sun.

Several results have been obtained.

First. The action increases with the density of the air, and

is diminished as it becomes more rarified.

The experiments were made with an air-pump and two cylindrical receivers of the same size, about four inches in diameter and thirty in length. In each were placed two thermometers, and the air was exhausted from one and condensed in the other. After both had acquired the same temperature they were placed in the sun, side by side, and while the action of the sun's rays rose to 110° in the condensed tube, it attained only 88° in the other. I had no means at hand of measuring the degree of condensation or rarefaction.

The observations taken once in two or three minutes, were as follows:

Eshan	ted Tube	Condensed Tube.						
In shade.	In eas.	In chade,	In son.					
75	80	76	80					
76	82	78	95					
80	81	80	100					
83	86	82	105					
84	88	85	110					

This circumstance must affect the power of the sun's rays in different places, and contribute to produce their feeble action on the summits of lofty mountains.

Secontly. The action of the sun's rays was found to be greater in moist than in dry air.

In one of the receivers the air was saturated with moisturein the other it was dried by the use of chlorid of calcium.

Both were placed in the sun as before and the result was as follows:

Dry	Als.	Damp Ale.						
In shade,	In sun.	In shade.	In ope.					
75	75	75	7.5					
18	88	. 78	90					
82	102	82 82	106					
81 82	104		110					
88	105	92	114					

The high temperature of moist air has frequently been observed. Who has not experienced the burning heat of the sun that precedes a summer's shower? The isothermal lines will, I think, be found to be much affected by the different degrees of moisture in different places.

Thirdly. The highest effect of the sun's rays I have found to be in carbonic acid gas.

One of the receivers was filled with it, the other with common air, and the result was as follows:

In Come	son Air.	In Curbosin	Arid Gus.
In shade,	In won.	In shade.	Is out.
80	90	80	90
81 -	94	84	100
80	- 99	84	110
81	100	85	120

The receiver containing the gas became itself much heated very sensibly more so than the other—and on being removed, it was many times as long in cooling.

An atmosphere of that gas would give to our earth a high temperature; and if as some suppose, at one period of its history the air had mixed with it a larger proportion than at present, an increased temperature from its own action as well as from increased weight must have necessarily resulted.

On comparing the sun's heat in different gases, I found it to be in hydrogen gas, 104°; in common air, 106°; in oxygen gas, 108°; and in carbonic acid gas, 125°.


ART. XXXII.—Review of a portion of the Geological Map of the United States and British Provinces by Jules Marcon;\* by WIL-MAK P. BLAKE.

Geological maps of the United States published in Europe and widely circulated among European geologists, are necessarily regarded by us with no small degree of attention and curiosity. This is more especially true, when such maps embrace regions of which the geography has only recently been made known and the geology has never before been laid down on a map with any approach to accuracy.

The recent geological map and profile by M. J. Marcou, which has appeared in the Annales des Mines and in the Bulletin of

\* Carte Géologique des Etats-Unis et des Provinces Anglaises de l'Amérique du Nard par Jules Marcosa. Annales des Mines, 2s Série, T. vii, p. 329. Published the min.


Résumé explicatif d'une carte géologique des Etate-Unis et des provinces auglaines de l'Amérique du Nord, avec un profit géologique allant de la vallée du Mutasippi aux côtes du Pacifique, et une planche de fessiles, par M. Jules Marcou Bulletin de la Société Géologique de Prance. Mai, 1855, p. 813.



1861, Tyndall's apparatus measuring properties of greenhouse gases.

John Tyndall FRS (2 August 1820 – 4 December 1893) was a prominent 19th-century Irish physicist. His scientific fame arose in the 1850s from his study of diamagnetism.

Later he made discoveries in the realms of infrared radiation and the physical properties of air, proving the connection between atmospheric CO2 and what is now known as the greenhouse effect in 1859.



#### ON RADIATION.

THE "REDE" LECTURE, DELIVERED IN THE SENATE-HOUSE, BEFORE THE UNIVERSITY OF CAMBRIDGE, ENGLAND, ON TUESDAY, MAY 16, 1865.

1.	Visible and Invisible Radiation	5
2.	Origin and Character of Radiation. The Ether	9
3.	The Atomic Theory in Reference to the Ether	12
	Absorption of Radiant Heat by Gases	
5.	Formation of Invisible Foci	17
6.	Visible and Invisible Rays of the Electric Light	19
7.	Combustion by Invisible Rays	21
8.	Transmutation of Rays. Calorescence	23
9.	Deadness of the Optic Nerve to the Calorific Rays	25
	Persistence of Rays	
11.	Absorption of Radiant Heat by Vapours and Odours	81
12.	Aqueous Vapour in Relation to Terrestrial Temperatures	88
13.	Liquids and their Vapours in Relation to Radiant Heat	36
14.	Reciprocity of Radiation and Absorption	37
15.	Influence of Vibrating Period and Molecular Form. Physical	
	Analysis of the Human Breath	40
16.	Summary and Conclusion	

"De Saussure, Fourier, M. Pouillet, and Mr. Hopkins regard this interception of the terrestrial rays as exercising the most important influence on climate. . . every variation [in aqueous vapour] must produce a change of climate. Similar remarks would apply to the carbonic acid diffused through the air, while an almost inappreciable admixture of any of the hydrocarbon vapours would produce great effects on the terrestrial rays and produce corresponding changes of climate. It is not, therefore, necessary to assume alterations in the density and height of the atmosphere to account for different amounts of heat being preserved to the earth at different times; a slight change in its variable constituents would suffice for this. Such changes in fact may have produced all the mutations of climate which the researches of geologists reveal."



# On the Influence of Carbonic Acid in the Air upon the Temperature of the Ground

Svante Arrhenius

Philosophical Magazine and Journal of Science Series 5, Volume 41, April 1896, pages 237-276.

This photocopy was prepared by Robert A. Rohde for Global Warming Art (http://www.globalwarmingart.com/) from original printed material that is now in the public domain.

Arrhenius's paper is the first to quantify the contribution of carbon dioxide to the greenhouse effect (Sections I-IV) and to speculate about whether variations in the atmospheric concentration of carbon dioxide have contributed to long-term variations in climate (Section V). Throughout this paper, Arrhenius refers to carbon dioxide as "carbonic acid" in accordance with the convention at the time he was writing.

Contrary to some misunderstandings, Arrhenius does not explicitly suggest in this paper that the burning of fossil fuels will cause global warming, though it is clear that he is aware that fossil fuels are a potentially significant source of carbon dioxide (page 270), and he does explicitly suggest this outcome in later work.

LONDON, EDINBURGH, AND DUBLIN

# PHILOSOPHICAL MAGAZINE

JOURNAL OF SCIENCE.

FIFTH SERIES.]

APRIL 1896.

XXXI. On the Influence of Carbonic Acid in the Air upon the Temperature of the Ground. By Prof. Syante ARRHENIUS \*.

> 1. Introduction: Observations of Langley on Atmospherical Absorption.

GREAT deal has been written on the influence of A the absorption of the atmosphere upon the climate. Tyndail † in particular has pointed out the enormous importance of this question. To him it was chiefly the diurnal and annual variations of the temperature that were lessened by this circumstance. Another side of the question, that has long attracted the attention of physicists, is this: Is the mean temperature of the ground in any way influenced by the presence of heat-absorbing gases in the atmosphere? Fourier: maintained that the atmosphere acts like the glass of a hothouse, because it lets through the light rays of the sun but retains the dark rays from the ground. This idea was elaborated by Pouillet §; and Langley was by some of his researches led to the view, that "the temperature of the earth under direct sunshine, even though our atmosphere were present as now, would probably fall to -200° C., if that atmosphere did not possess the quality of selective

Mem. de l'Ac. R. d. Sci. de l'Inst. de France, t. vii. 1827.

Comptes rendus, t, vii. p. 41 (1838).

Phil Man S 5 Val 41 No 051 4 7 1900

Extract from a paper presented to the Royal Swedish Academy of Sciences, 11th December, 1895. Communicated by the Author.
 † 'Heat a Mode of Motion,' 2nd ed. p. 405 (Lond., 1865).

Table VII .- Variation of Temperature caused by a given Variation of Carbonic Acid.

de.		Carbonic Acid=067.					Carbonic Acid=1.5.						Carbonic Acid=2.0.						id=2	Carbonic Acid=3.0.					
Latitude.	Dec Feb.	March- May.	June- Aug.	Sept Nov.	Mean of the year.	Dec Feb.	March- May.	June- Aug.	Sept Nov.	Mean of the year.	Dec Feb.	March- May.	June- Aug.	Sept Nov.	Mean of the year.	Dec Feb.	March- May.	June-	Sept Nov.	Mean of the year.	Dec Feb.	March- May.	June-	Sept Nov.	Mean of
70	-2.9	-3.0	-3.4	-3.1	-3.1	3.3	3.4	3.8	3.6	3.52	6:0	6.1	6.0	6.1	6:05	7.9	8.0	7:9	8.0	7:95	9-1	9:3	9.4	9.4	9:
60	1		33.		2000	00000			100000	3.62	STANDED !	6.1	1000		6-02		8:0	7.6	7-9	7.87	9-3	9.5	8.9	9.5	9-
50	-3.0	-3.2	-3.4	-3.3	-3.22	3.4	3.7	3.6	3.8		6.1		5.8	6.1		8.0				1.01			-		
40	-3.2	-3.3	-3:3	-34	-3.3	3.7	3.8	3.4	3.7	3.65	6.1	6.1	5.5	6.0	5.92	8-0	7.9	7.0	7.9	7.7	9.5	9.4	8.6	9.2	9.
	-3.4	-3.4	-3.2	-3.3	-3.32	3.7	3.6	3.3	3.5	3.52	6.0	5.8	5.4	5.6	5.7	7.9	7.6	6.9	7.3	7.42	9.3	9.0	8.2	8.8	8
30	-3.3	-3.2	-3.1	-3.1	-3.17	3.5	3.3	3.2	3.5	3.47	56	5.4	50	5.2	5.3	7.2	7.0	6.6	6.7	6.87	8.7	8.3	7.5	7.9	8
20	-3.1	-3.1	-3.0	-3.1	-3:07	3.5	3.2	3.1	3.2	3.25	5.2	5.0	4.9	5.0	5.02	6.7	6.6	6.3	6.6	6.52	7.9	7.5	7.2	7.5	7:
10	-3.1	-3.0	-30	-3.0	-3.02	3.2	3.2	3.1	3.1	3.15	5.0	5.0	4.9	4.9	4.95	6.6	6.4	6.3	6.4	6.42	7.4	7:3	7.2	7.3	7
0	-30	-3.0	-3.1	-3.0	-3.02	3.1	3.1	3.2	3.2	3:15	100000	4.0		5.0	4.95	6-4	6.4	6-6	6-6	6.5	7.3	7.3	7.4	7:4	7
-10													5-0			07.57		100000	03000			0.55	1000000	1000	
-20	-3.1	-3.1	-3.2	-3.1	-3.12	3.2	3.2	3.2	3.2	3.2	5.0	5.0	5.2	5.1	5.07	6-6	6.6	6.7	6.7	6.65	7.4	7.5	8.0		1
-30	-3.1	-3.2	-3.3	-3.2	-3.2	3.2	3.2	3.4	3.3	3.27	5.2	5.3	5.5	5.4	5.35	6.7	6.8	7.0	7.0	6.87	7.9	8.1	8.6	8.3	8
	-3.3	-3.3	-3.4	-3.4	-3.35	3.4	3.5	3.7	3.5	3.52	5.5	56	5.8	5.6	5.62	7.0	7.2	7.7	7.4	7.32	8.6	8.7	9.1	8.8	8
-40	-3.4	-3.4	-3.3	-3.4	-3:37	3.6	3.7	3.8	37	3.7	5.8	6-0	60	6.0	5.95	77	7.9	7.9	7.9	7.85	9.1	9.2	9.4	9.3	9
-50 -60	-3.2	- 3.3	_	_	_	3.8	1		_	-	6.0	6-1	-	-	_	7.9	8.0	_	_	-	9.4	9.5	-		-

#### ADDENDUM\*.

As the nebulosity is very different in different latitudes, and also different over the sea and over the continents, it is evident that the influence of a variation in the carbonic acid of the air will be somewhat different from that calculated above, where it is assumed that the nebulosity is the same over the whole globe. I have therefore estimated the nebulosity at different latitudes with the help of the chart published by Teisserenc de Bort, and calculated the following table for

e.	Nebulo	sity.	11	Reduc	tion fac	tor.	K=0	67.	K=1.5.		
Latitude.	Conti- nent.	Осезп.	Continent per cent.	Conti-	Oeean.	Mean.	Conti- nent.	Ocean.	Conti- nent.	Ocean,	
70	58:1	66.7	72:1	0.899	0.775	0.864	-2.8	-24	3·1	2.7	
60	56:3	67-6	55-8	0.924	0.763	0.853	-30	-2.4	3.3	2.7	
50	45.7	63:3	529	1.057	0.813	0.942	-3.5	-27	38	2.9	
40	36-5	52.5	429	1.177	0 939	1-041	-39	-3.1	4.1	3.3	
30	28-5	47-2	38-8	1.296	1.009	1.120	-41	-3.2	4.5	3-1	
20	28-5	47:0	24-2	1.308	1.017	1.087	-4:1	-3.2	4.3	3.	
10	50.1	56.7	23.3	1.031	0.903	0.933	- 3.1	-2.7	3.3	24	
0	54.8	59-7	24-2	0.97	0.867	0.892	-2.9	-26	3.1	24	
-10	47.8	54:0	22.5	1.056	0.932	0.96	-3.3	-29	3.4	34	
-20	29-6	49-6	23-3	1-279	0.979	0.972	-4.1	-3.1	4.2	3	
-30	38.9	51-0	12-5	1:152	0.958	0.982	-3.8	-3.2	4.0	3	
-40	62.0	61:1	2-5	0.86	0.837	0.838	-29	-2.8	,3.2	3	
-50 -60	71-0	71.5	0.9	0.749	0.719	0.719					

\* Cf. p. 265.

Arrhenius estimated also cooling effect of atmospheric aerosols!


#### An Attempt to Frame a Working Hypothesis of the Cause of Glacial Periods on an Atmospheric Basis

### T. C. Chamberlin The Journal of Geology Vol. 7, No. 6 (Sep. - Oct., 1899), pp. 545-584

#### AN ATTEMPT TO FRAME A WORKING HYPOTHESIS OF THE CAUSE OF GLACIAL PERIODS ON AN ATMOSPHERIC BASIS'

THERE are hypotheses and working hypotheses. The suggestion that the last glacial period was caused by the passage of the solar system through a cold region of space may be styled a hypothesis, but scarcely a working hypothesis in the geological sense, for it does not form the groundwork or incentive of geological inquiry. An astronomer might be moved to hunt for the cold spot, but it has no inspiration for the geologist. General suggestions of a possible cause do not reach the dignity of working hypotheses until they are given concrete form, are fitted in detail to the specific phenomena, and are made the agents of calling into play effective lines of research. The construction of a concrete working hypothesis suited to stimulate and guide investigation in a wholesome manner, and to take its place in competition with other hypotheses of like working potentialities, thereby inducing a more searching scrutiny of the phenomena and a more varied application of interpretations, represents the higher limit of present reasonable aspiration. It is much too ambitious to hope for a demonstrative solution of the origin of the earth's glacial periods by first intention in the present state of knowledge.


The hypothesis here offered is not worked out into satisfactory detail at all points, but it is hoped that it is sufficiently matured to justify a preliminary statement. In forming it, which has been the work of several years, I have found, or seemed to find, the phenomena of past glaciation intimately associated with a long chain of other phenomena to which at




A brief statement of the salient features of this hypothesis was given in a paper entitled A Group of Hypotheses Bearing on Climatic Changes, Jour. Geol., Vol. V., pp. 653-683, Oct.-Nov. 1897. For earlier history see footnotes on pp. 654 and 681 of that paper.



# Milankovitch Cycles




#### Milankovitch Cycles and Temperature from Vostok Ice-core



# QUARTERLY JOURNAL ROYAL METEOROLOGICAL SOCIETY





#### The artificial production of carbon dioxide and its influence on temperature

G. S. Callendar Issue

Article first published online: 10 SEP 2007

DOI: 10.1002/qj.49706427503

Copyright @ 1938 Royal Meteorological Society



Quarterly Journal of the

Volume 64, Issue 275, pages 223-240. April 1938

Royal Meteorological Society





Additional Information (Show All)

How to Cite Author Information Publication History

Abstract

Article

References

Cited By



Cet PDF (1012K)

#### Abstract

By fuel combustion man has added about 150,000 million tons of carbon dioxide to the air during the past half century. The author estimates from the best available data that approximately three quarters of this has remained in the atmosphere.

The radiation absorption coefficients of carbon dioxide and water vapour are used to show the effect of carbon dioxide on "sky radiation." From this the increase in mean temperature, due to the artificial production of carbon dioxide, is estimated to be at the rate of 0.003°C, per year at the present time.

The temperature observations at zoo meteorological stations are used to show that world temperatures have actually increased at an average rate of 0.005°C, per year during the past half century.



- 1956 -Phillips produces a computer model of the global atmosphere, Plass calculates again that radiation balance depends on CO<sub>2</sub> concentration
- 1957 -Revelle and Suess demonstrate that only a fraction (about ½) of CO<sub>2</sub> produced by humans absorbed by the oceans.
- 1960 Keeling detects an annual rise of CO<sub>2</sub> in the atmosphere.
- 1965 Lorenz points out the chaotic nature of climate system and the possibility of sudden shifts.
- 1966 Emiliani's analysis of deep-sea cores and Broecker's analysis of corals show that the timing of ice ages was set by small orbital shifts.
- **1979** US National Academy of Sciences report finds that doubling CO<sub>2</sub> will bring 1.5-4.5°C global warming.
- 1990 First IPCC report says world has been warming and future warming seems likely.
- 1991 Mt. Pinatubo explodes; Hansen predicts cooling pattern, verifying (by 1995) computer models of aerosol effects.

# Carbon Dioxide Exchange Between Atmosphere and Ocean and the Question of an Increase of Atmospheric CO, during the Past Decades

By ROGER REVELLE and HANS E. SUESS, Scripps Institution of Oceanography, University of California, La Jolla, California

(Manuscript received September 4, 1956)

#### Abstract

From a comparison of C<sup>14</sup>/C<sup>18</sup> and C<sup>13</sup>/C<sup>18</sup> ratios in wood and in marine material and from a slight decrease of the C<sup>14</sup> concentration in terrestrial plants over the past 50 years it can be concluded that the average lifetime of a CO<sub>4</sub> molecule in the atmosphere before it is dissolved into the sea is of the order of 10 years. This means that most of the CO<sub>4</sub> released by artificial fuel combustion since the beginning of the industrial revolution must have been absorbed by the oceans. The increase of atmospheric CO<sub>4</sub> from this cause is at present small but may become significant during future decades if industrial fuel combustion continues to rise exponentially.

Present data on the total amount of  $CO_2$  in the atmosphere, on the rates and mechanisms of exchange, and on possible fluctuations in terrestrial and marine organic carbon, are inadequate for accurate measurement of future changes in atmospheric  $CO_1$ . An opportunity exists during the International Geophysical Year to obtain much of the necessary information.

(Tellus, 9, 1957, 18–27)

"Human beings are now carrying out a large scale geophysical experiment of a kind that could not have happened in the past nor be reproduced in the future. Within a few centuries, we are returning to the atmosphere and oceans the concentrated organic carbon stored in sedimentary rocks over hundreds of millions of years."





# The Carbon Dioxide Theory of Climatic Change

By GILBERT N. PLASS

The Johns Hopkins University, Baltimore, Md.1

(Manuscript received August 9 1955)

## Abstract

The most recent calculations of the infra-red flux in the region of the 15 micron CO2 band show that the average surface temperature of the earth increases 3.6° C if the CO2 concentration in the atmosphere is doubled and decreases 3.8° C if the CO2 amount is halved, provided that no other factors change which influence the radiation balance. Variations in CO, amount of this magnitude must have occurred during geological history; the resulting temperature changes were sufficiently large to influence the climate. The CO2 balance is discussed. The CO2 equilibrium between atmosphere and oceans is calculated with and without CaCO3 equilibrium, assuming that the average temperature changes with the CO2 concentration by the amount predicted by the CO2 theory. When the total CO2 is reduced below a critical value, it is found that the climate continuously oscillates between a glacial and an inter-glacial stage with a period of tens of thousands of years; there is no possible stable state for the climate. Simple explanations are provided by the CO2 theory for the increased precipitation at the onset of a glacial period, the time lag of millions of years between periods of mountain building and the ensuing glaciation, and the severe glaciation at the end of the Carboniferous. The extra CO2 released into the atmosphere by industrial processes and other human activities may have caused the temperature rise during the present century. In contrast with other theories of climate, the CO2 theory predicts that this warming trend will continue, at least for several centuries.

Introduction may have produced all the mutations of

# Wallace S. Broecker, Science, Vol. 189, No. 4201 (Aug. 8, 1975), 460-463

# Climatic Change: Are We on the Brink of a

## **Pronounced Global Warming?**

Abstract. If man-made dust is unimportant as a major cause of climatic change, then a strong case can be made that the present cooling trend will, within a decade or so, give way to a pronounced warming induced by carbon dioxide. By analogy with similar events in the past, the natural climatic cooling which, since 1940, has more than compensated for the carbon dioxide effect, will soon bottom out. Once this happens, the exponential rise in the atmospheric carbon dioxide content will tend to become a significant factor and by early in the next century will have driven the mean planetary temperature beyond the limits experienced during the last 1000 years.





Fig. 1. Curves for the global temperature change due to chemical fuel CO<sub>2</sub>, natural climatic cycles, and the sum of the two effects. The measured temperature anomaly for successive 5-year means from meteorological records over the last century is given for comparison.



# SCIENCE

# Climate Impact of Increasing Atmospheric Carbon Dioxide

J. Hansen, D. Johnson, A. Lacis, S. Lebedeff
P. Lee, D. Rind, G. Russell





Fig. 6. Projections of global temperature. The diffusion coefficient beneath the ocean mixed layer is 1.2 cm<sup>2</sup> sec<sup>-1</sup>, as required for best fit of the model and observations for the period 1880 to 1978. Estimated global mean warming in earlier warm periods is indicated on the right.






# The Nobel Prize in Physics 2021



III. Niklas Elmehed © Nobel Prize Outreach

Syukuro Manabe

Prize share: 1/4



III. Niklas Elmehed © Nobel Prize Outreach

Klaus Hasselmann

Prize share: 1/4



III. Niklas Elmehed © Nobel Prize Outreach

Giorgio Parisi

Prize share: 1/2

The Nobel Prize in Physics 2021 was awarded "for groundbreaking contributions to our understanding of complex systems" with one half jointly to Syukuro Manabe and Klaus Hasselmann "for the physical modelling of Earth's climate, quantifying variability and reliably predicting global warming" and the other half to Giorgio Parisi "for the discovery of the interplay of disorder and fluctuations in physical systems from atomic to planetary scales."

# Thermal Equilibrium of the Atmosphere with a Convective Adjustment

#### SYUKURO MANABE AND ROBERT F. STRICKLER

General Circulation Research Laboratory, U. S. Weather Bureau, Washington, D. C. (Manuscript received 19 December 1963, in revised form 13 April 1964)




Fig. 6c. Thermal equilibrium of various atmospheres which have a critical lapse rate of 6.5 deg km<sup>-1</sup>. Vertical distributions of gaseous absorbers at 35N, April, were used.  $S_c=2$  ly min<sup>-1</sup>,  $\cos \bar{f} = 0.5$ , r = 0.5, no clouds.




Fig. 4. The dashed, dotted, and solid lines show the thermal equilibrium with a critical lapse rate of 6.5 deg km<sup>-1</sup>, a dryadiabatic critical lapse rate (10 deg km<sup>-1</sup>), and pure radiative equilibrium.




Fig. 8. Vertical distribution of radiative convective equilibrium temperature of the atmosphere with a given distribution of relative humidity for various values of the solar constant.




Fig. 12. Vertical distributions of radiative convective equilibrium temperature for various values of water vapor mixing ratio in the stratosphere.

## Stochastic climate models

## Part I. Theory

By K. HASSELMANN, Max-Planck-Institut für Meteorologie, Hamburg, FRG

(Manuscript received January 19; in final form April 5, 1976)

#### ABSTRACT

A stochastic model of climate variability is considered in which slow changes of climate are explained as the integral response to continuous random excitation by short period "weather" disturbances. The coupled ocean-atmosphere-cryosphere-land system is divided into a rapidly varying "weather" system (essentially the atmosphere) and a slowly responding "climate" system (the ocean, cryosphere, land vegetation, etc.). In the usual Statistical Dynamical Model (SDM) only the average transport effects of the rapidly varying weather components are parameterised in the climate system. The

arise only through variable external conditions. The essential feature of stochastic climate models is that the non-averaged "weather" components are also retained. They appear formally as random forcing terms. The climate system, acting as an integrator of this short-period excitation, exhibits the same random-walk response characteristics as large particles interacting with an ensemble of much smaller particles in the analogous Brownian motion problem. The model predicts "red" variance spectra, in qualitative agreement with observations. The evolution of the climate probability distribution is described by a Fokker-Planck equation, in which the effect of the random weather excitation is represented by diffusion terms. Without stabilising feedback, the model predicts a continuous increase in climate variability, in analogy with the continuous, unbounded dispersion of particles in Brownian motion (or in a homogeneous turbulent fluid). Stabilising feedback yields a statistically stationary climate probability distribution. Feedback also results in a finite degree of climate predictability, but for a stationary climate the predictability is limited to maximal skill parameters of order 0.5.

#### REFERENCES

- Budyko, M. I. 1969. The effect of solar radiation variations on the climate of the earth. Tellus 21. 611-619
- Frankignoul, C. & Hasselmann, K. 1976, Stochastic elimate models. Part 2, Application to sea-surface temperature anomalies and thermocline variability (in preparation).
- GARP US Committee Report, 1975. Understanding climate change. A programme for action. Nat. Acad. Sciences, Wash.
- GARP Publication 16, 1975. The physical basis of climate and climate modelling, World Met. Organiz., Internat. Council Scient. Unions.
- Hasselmann, K. 1966. Feynman diagrams and interaction rules of wave-wave scattering processes. Rev. Geophys. 4, 1-32.
- Hasselmann, K. 1967, Non-linear interactions treated by the methods of theoretical physics (with application to the generation of waves by wind). Proc. Roy. Soc. A 299, 77-100.
- Hinze, J. O. 1959. Turbulence, McGraw-Hill.
- Kadomtsev, B. B. 1965. Plasma turbulence, Academic Press.
- King, J. W. 1975. Sun-weather-relationships. Aeronautics and Astronautics 13, 10-19.
- Lemke, P. 1976. Stochastic climate models, Part 3, Application to zonally averaged energy models (in preparation).

- Lorenz, E. N. 1965, A study of the predictability of a 28-variable atmospheric model. Tellus 17, 321 - 333.
- Lorenz, E. N. 1968, Climate determinism. Meleor. Monographs 8, 1-3,
  - sea interaction and climatic fluctuation. (Symp. on the Arctic Heat Budget and Atmospheric Circulation, Lake Arrowhead, Calif., 1966.) Mem. RM-5233-NSF, The Rand Corp., Santa Monica.
- Monin, A. S. & Vulis, I. L. 1971. On the spectra of long-period oscillations of geophysical parameters. Tellus 23, 337-345.
- Sellers, W. D. 1969. A global climate model based on the energy balance of the earth-atmosphere system. J. Appl. Met, 8, 392-400.
- Tatarski, V. I. 1961. Wave propagation in a turbulent medium, McGraw-Hill.
- Taylor, G. I. 1921. Diffusion by continuous movements, Proc. Lond. Math. Soc. 20, 196.
- Wang, M. C. & Uhlenbek, G. E. 1945. On the theory of the Brownian motion. Rev. Mod. Phys. 17, 323 - 342.
- Wilcox, J. M. 1975. Solar activity and the weather. J. Atmosph. Terrestr. Phys. 37, 237-256.

## Stochastic resonance in climatic change

By ROBERTO BENZI, Istituto di Fisica dell'Atmosfera, C.N.R., Piazza Luizi Sturzo 31, 00144, Roma,

GIORGIO PARISI, I.N.F.N., Laboratori Naziona Francati, Francati, Roma, Italy,

Ann. Hartford, Connecticut 06120, U.S.A.

and ANGELO VULPIANI, Istituto di Fisica "G. Marconi", Università di Roma, Italy

(Manuscriet received November 12, 1980; in final form March 13, 1981)

#### ABSTRACT

An amplification of random perturbations by the interaction of non-linearities internal to the climatic system with external, orbital forcing is found. This stochastic resonance is investigated in a highly simplified, pero-dimensional climate model. It is conceivable that this new type of resonance might play a role in explaining the 10° year peak in the power spectra of pulsoclimatic records.

B. BENZI ET AL

#### REFERENCES

Blattacharya, K. and Glol, M. 1978. An energy belance: model with multiply periodic and quasi-chaotic free oscillations. In Evolution of planetary atmospheres and climateless of the earth. Toolsway, France: Nicolia, C. 1980. Pluctuations, solar periodicities and Centre National d'Enades Speciales, 299-310.

Budyles, M. L. 1969. The offers of solar radiation.

Ghil, M. 1976. Climate stability for a Sellers-type model. J. Almon. Sci. JJ, 3-20.

Chil, M. 1980. Internal climatic mechanisms particlputing in glaciations cycles. In Climate participes and cartability (ed. A. Berger). Doedrocki Boston-London: D. Reidel Publ. Co., in press.

Ghil, M. and Bhattacharya, K. 1979. An energy bulance model of glaciation cycles. In Climate models (ed. W. L. Gatest, WMO/GCSU, Geneva, Switzerland: GARP Publ. Series, n. 22, 886-916.

Gibran, I. I. and Skorobod, A. V. 1972. Stochastic differential equations. Berlin-Heidelberg-New York:

Hasselman, K. 1976. Stockastic climate models, part II. Theory, Feller 28, 423-484.

Variations in the earth's orbit: pagemaker of the icenges, Science 194, 1121-1112.

Milankovitch, M. 1930. Handbuck der Klimanningte F. Tell lads, A. W. Koppen and W. Wegener). Serlin: Doppen and Geiger:

climatic transitions. In Proc. Int. Conf. Sun and Climate, CNES, Toulouse.

variations on the climate of the earth. Tolks 21. Nicolis, G. and Nicolis, C. 1981. Suchastic aspects of climatic transitions—additive fluctuations. Tellus III.

> North, G. R. and Coakly, Jr. J. A. 1979. Differences between seasonal and mean annual energy belance model calculations of climate and climate sensitivity. J. Jones, Sci. M. 1189-1204.

> Poland, D., Ingmoli, A. P. and Lockwood, J. G. 1980. Response of a sonal climate-tenhest model to the critical perturbations during the quaternary ice ages. Tellio JC, 301-319.

> Schoolder, S. H. and Thompson, S. L. 1979. Its ages and orbital variations; some simple theory and modelling. Quat. Res. 12, 168-203.

ear, M. J. and Held, I. M. 1979. The sometimes of an vergy balance climate model to variations in the rbital parameters. J. Geophys. Res. 84, 4825-4836. ra. A. 1981. On southeric perturbation and longterm climate behaviour, Opart, J. Roy, Meteoroi, Soc.

JULY 137-152.

# 5. Conclusions

Our results point to the possibility of explaining large amplitude, long-term alternations of temperature by means of a co-operation between external periodic forcing due to orbital variations and an internal stochastic mechanism. The external periodic forcing alone is unable to reproduce the major peak in the observed quaternary climate records. The internal stochastic forcing alone does not reproduce it either. The combination of the two effects, however, produces what we may call a stochastic resonance, which amplifies the small external forcing: a small change in the external forcing induces a large change in the probability of jumping between two observable climates. This new mechanism could be useful in our understanding of long-term climatic change. At any rate, it seems to warrant further investigation.

# Carbon Dioxide and Climate: A Scientific Assessment

# Ad Hoc Study Group on Carbon Dioxide and Climate

Report of an Ad Hoc Study Group on Carbon Dioxide and Climate Woods Hole, Massachusetts July 23-27, 1979 to the Climate Research Board Assembly of Mathematical and Physical Sciences National Research Council



Jule G. Charney, Massachusetts Institute of Technology, Chairman Akio Arakawa, University of California, Los Angeles D. James Baker, University of Washington Bert Bolin, University of Stockholm Robert E. Dickinson, National Center for Atmospheric Research Richard M. Goody, Harvard University Cecil E. Leith, National Center for Atmospheric Research Henry M. Stommel, Woods Hole Oceanographic Institution Carl I. Wunsch, Massachusetts Institute of Technology We have examined the principal attempts to simulate the effects of increased atmospheric CO<sub>2</sub> on climate. In doing so, we have limited our considerations to the direct climatic effects of steadily rising atmospheric concentrations of CO<sub>2</sub> and have assumed a rate of CO<sub>2</sub> increase that would lead to a doubling of airborne concentrations by some time in the first half of the twenty-first century.


When it is assumed that the CO<sub>2</sub> content of the atmosphere is doubled and statistical thermal equilibrium is achieved, the more realistic of the modeling efforts predict a global surface warming of between 2°C and 3.5°C, with greater increases at high latitudes. This range reflects both uncertainties in physical understanding and inaccuracies arising from the need to reduce the mathematical problem to one that can be handled by even the fastest available electronic computers. It is significant, however, that none of the model calculations predicts negligible warming.

The primary effect of an increase of CO<sub>2</sub> is to cause more absorption of thermal radiation from the earth's surface and thus to increase the air temperature in the troposphere. A strong positive feedback mechanism is the accompanying increase of moisture, which is an even more powerful absorber

of terrestrial radiation. We have examined with care all known negative feed-back mechanisms, such as increase in low or middle cloud amount, and have concluded that the oversimplifications and inaccuracies in the models are not likely to have vitiated the principal conclusion that there will be appreciable warming. The known negative feedback mechanisms can reduce the warming, but they do not appear to be so strong as the positive moisture feedback. We estimate the most probable global warming for a doubling of CO<sub>2</sub> to be near 3°C with a probable error of ± 1.5°C. Our estimate is based primarily on our review of a series of calculations with three-dimensional models of the global atmospheric circulation, which is summarized in Chapter 4. We have also reviewed simpler models that appear to contain the main physical factors. These give qualitatively similar results.

To summarize, we have tried but have been unable to find any overlooked or underestimated physical effects that could reduce the currently estimated global warmings due to a doubling of atmospheric CO<sub>2</sub> to negligible proportions or reverse them altogether. However, we believe it quite possible that the capacity of the intermediate waters of the oceans to absorb heat could delay the estimated warming by several decades. It appears that the warming will eventually occur, and the associated regional climatic changes so important to the assessment of socioeconomic consequences may well be significant, but unfortunately the latter cannot yet be adequately projected.


The Intergovernmental Panel on Climate Change (IPCC), established in 1988 is the United Nations body for assessing the science related to climate change




## IPCC First Assessment Report 1990 (FAR)


(OUT OF PRINT) Digitized by the Digitization and Microform Unit, UNOG Library, 2010 (Available in English only except where stated)







Working Group II: Impacts Assessment of Climate Change



Working Group III: The IPCC Response Strategies

First Assessment Report Overview Chapter (PDF)

# 1.0.1 We are certain of the following:

- There is a natural greenhouse effect which already
   keeps the Earth warmer than it would otherwise be.
- Emissions resulting from human activities are substantially increasing the atmospheric concentrations of the greenhouse gases: carbon dioxide, methane, chlorofluorocarbons (CFCs) and nitrous oxide. These increases will enhance the greenhouse effect, resulting on average in an additional warming of the Earth's surface. The main greenhouse gas, water vapour, will increase in response to global warming and further enhance it.

# 1.0.3 Based on current model results, we predict:

An average rate of increase of global mean temperature during the next century of about 0.3°C per decade (with an uncertainty range of 0.2—0.5°C per decade) assuming the IPCC Scenario A (Business-as-Usual) emissions of greenhouse gases; this is a more rapid increase than seen over the past 10,000 years. This will result in a likely increase in the global mean temperature of about 1°C above the present value by 2025 (about 2°C above that in the pre-industrial period), and 3°C above today's value before the end of the next century (about 4°C above pre-industrial). The rise will not be steady because of other factors.

# 2. Impacts

- 2.0.1 The report on impacts of Working Group II is based on the work of a number of subgroups, using independent studies which have used different methodologies. Based on the existing literature, the studies have used several scenarios to assess the potential impacts of climate change. These have the features of:
- i) an effective doubling of CO<sub>2</sub> in the atmosphere between now and 2025 to 2050;
- ii) a consequent increase of global mean temperature in the range of 1.5°C to 4.5°C;
- iii) an unequal global distribution of this temperature increase, namely a smaller increase of half the global mean in the tropical regions and a larger increase of twice the global mean in the polar regions;
- iv) a sea-level rise of about 0.3—0.5 m by 2050 and about 1 m by 2100, together with a rise in the temperature of the surface ocean layer of between 0.2° and 2.5°C.

## 1992 Supplementary Reports

(OUT OF PRINT) Digitized by the Digitization and Microform Unit, UNOG Library, 2010 (Available in English only except where stated)



The Supplementary Report to The IPCC Scientific Assessment



The Supplementary Report to The IPCC Impacts Assessment



The IPCC 1990 and 1992 Assessments


CLICK HERE

CLICK HERE

CLICK HERE

#### IPCC Second Assessment Report: Climate Change 1995 (SAR)

(OUT OF PRINT) Digitized by the Digitization and Microform Unit, UNOG Library, 2010. (Available in English only except where stated)



Working Group I: The Science of Climate Change Full Report (PDF)



Working Group II: Impacts, Adaptations and Mitigation of Climate Change: Scientific-Technical Analyses Full Report (PDF)



Working Group III: Economic and Social Dimensions of Climate Change Full Report (PDF)




IPCC Second Assessment Full Report (PDF) Errata

Arabic - Chinese - French-Russian - Spanish


## IPCC Third Assessment Report: Climate Change 2001 (TAR)




Working Group I: The Scientific Basis



Working Group II: Impacts, Adaptation and Vulnerability



Working Group III: Mitigation



Synthesis Report

#### WG1 - Summary for Policymakers

The Third Assessment Report of Working Group I of the Intergovernmental Panel on Climate Change (IPCC) builds upon past assessments and incorporates new results from the past five years of research on climate change!. Many hundreds of scientists? from many countries participated in its preparation and review.


This Summary for Policymakers (SPM), which was approved by IPCC member governments in Shanghai in January 20013, describes the current state of understanding of the climate system and provides estimates of its projected future evolution and their uncertainties. Further details can be found in the underlying report, and the appended Source Information provides cross references to the report's chapters.

#### An increasing body of observations gives a collective picture of a warming world and other changes in the climate system.

Since the release of the Second Assessment Report (SAR4), additional data from new studies of current and palaeoclimates, improved analysis of data sets, more rigorous evaluation of their quality, and comparisons among data from different sources have led to greater understanding of climate change.

#### The global average surface temperature has increased over the 20th century by about 0.6°C.

- The global average surface temperature (the average of near surface air temperature over land, and sea surface temperature) has increased since 1861.
  Over the 20th century the increase has been 0.6 ± 0.2°C<sub>2.5</sub>° (Figure 1a). This value is about 0.15°C larger than that estimated by the SAR for the period up to 1994, owing to the relatively high temperatures of the additional years (1995 to 2000) and improved methods of processing the data. These numbers take into account various adjustments, including urban heat island effects. The record shows a great deal of variability; for example, most of the warming occurred during the 20th century, during two periods, 1910 to 1945 and 1976 to 2000.
- . Globally, it is very likely that the 1990s was the warmest decade and 1998 the warmest year in the instrumental record, since 1861 (see Figure 1a).
- New analyses of proxy data for the Northern Hemisphere indicate that the increase in temperature in the 20th century is likely! to have been the largest of
  any century during the past 1,000 years. It is also likely! that, in the Northern Hemisphere, the 1990s was the warmest decade and 1998 the warmest year
  (Figure 1b). Because less data are available, less is known about annual averages prior to 1,000 years before present and for conditions prevailing in most
  of the Southern Hemisphere prior to 1861.
- On average, between 1950 and 1993, night-time daily minimum air temperatures over land increased by about 0.2\*C per decade. This is about twice the
  rate of increase in daytime daily maximum air temperatures (0.1\*C per decade). This has lengthened the freeze-free season in many mid- and high latitude
  regions. The increase in sea surface temperature over this period is about half that of the mean land surface air temperature.







Figure 1: Variations of the Earth's surface temperature over the last 140 years and the last millennium.

(a) The Earth's surface temperature is shown year by year (red bars) and approximately decade by decade (black line, a filtered annual curve suppressing fluctuations below near decadal


## IPCC Fourth Assessment Report: Climate Change 2007 (AR4)



Working Group I Report "The Physical Science Basis"



Working Group II Report "Impacts, Adaptation and Vulnerability"



Working Group III Report "Mitigation of Climate Change"




The AR4 Synthesis Report

# NOBEL PEACE PRIZE 2007 Intergovernmental Panel on Climate Change (IPCC) and Albert Amold Gore Jr.

# 1. Observed changes in climate and their effects

Warming of the climate system is unequivocal, as is now evident from observations of increases in global average air and ocean temperatures, widespread melting of snow and ice and rising global average sea level (Figure SPM.1). {1.1}



# 2. Causes of change

Changes in atmospheric concentrations of greenhouse gases (GHGs) and aerosols, land cover and solar radiation alter the energy balance of the climate system. [2.2]

Global GHG emissions due to human activities have grown since pre-industrial times, with an increase of 70% between 1970 and 2004 (Figure SPM.3).<sup>5</sup> {2.1}

Carbon dioxide (CO<sub>2</sub>) is the most important anthropogenic GHG. Its annual emissions grew by about 80% between 1970 and 2004. The long-term trend of declining CO<sub>2</sub> emissions per unit of energy supplied reversed after 2000. [2.1]

Global atmospheric concentrations of CO<sub>2</sub>, methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) have increased markedly as a result of human activities since 1750 and now far exceed pre-industrial values determined from ice cores spanning many thousands of years. {2.2}




Figure SPM.3. (a) Global annual emissions of anthropogenic GHGs from 1970 to 2004.<sup>5</sup> (b) Share of different anthropogenic GHGs in total emissions in 2004 in terms of carbon dioxide equivalents (CO<sub>2</sub>-eq). (c) Share of different sectors in total anthropogenic GHG emissions in 2004 in terms of CO<sub>2</sub>-eq. (Forestry includes deforestation.) (Figure 2.1)

## IPCC Fifth Assessment Report



Working Group I Report "Climate Change 2013: The Physical Science Basis"



Working Group II Report "Climate Change 2014: Impacts, Adaptation, and



Working Group III Report "Climate Change 2014: Mitigation of Climate Change"



"Climate Change 2014: Synthesis Report"


## SPM 1. Observed Changes and their Causes

Human influence on the climate system is clear, and recent anthropogenic emissions of greenhouse gases are the highest in history. Recent climate changes have had widespread impacts on human and natural systems. (1)


## SPM 1.1 Observed changes in the climate system

Warming of the climate system is unequivocal, and since the 1950s, many of the observed changes are unprecedented over decades to millennia. The atmosphere and ocean have warmed, the amounts of snow and ice have diminished, and sea level has risen. (1.1)

# https://www.ipcc.ch/reports/



https://www.ipcc.ch/report/ar6/syr/



A.1 Human activities, principally through emissions of greenhouse gases, have unequivocally caused global warming, with global surface temperature reaching 1.1°C above 1850-1900 in 2011-2020. Global greenhouse gas emissions have continued to increase, with unequal historical and ongoing contributions arising from unsustainable energy use, land use and land-use change, lifestyles and patterns of consumption and production across regions, between and within countries, and among individuals (high confidence). {2.1, Figure 2.1, Figure 2.2}

## A. The Current State of the Climate

- A.1 It is unequivocal that human influence has warmed the atmosphere, ocean and land.
  Widespread and rapid changes in the atmosphere, ocean, cryosphere and biosphere have occurred.
- A.2 The scale of recent changes across the climate system as a whole and the present state of many aspects of the climate system are unprecedented over many centuries to many thousands of years.

## B. Possible Climate Futures

- B.1 Global surface temperature will continue to increase until at least the mid-century under all emissions scenarios considered. Global warming of 1.5°C and 2°C will be exceeded during the 21st century unless deep reductions in carbon dioxide (CO<sub>2</sub>) and other greenhouse gas emissions occur in the coming decades.
- B.2 Many changes in the climate system become larger in direct relation to increasing global warming. They include increases in the frequency and intensity of hot extremes, marine heatwaves, and heavy precipitation, agricultural and ecological droughts in some regions, and proportion of intense tropical cyclones, as well as reductions in Arctic sea ice, snow cover and permafrost.


## C. Climate Information for Risk Assessment and Regional Adaptation

- C.1 Natural drivers and internal variability will modulate human-caused changes, especially at regional scales and in the near term, with little effect on centennial global warming. These modulations are important to consider in planning for the full range of possible changes.
- C.2 With further global warming, every region is projected to increasingly experience concurrent and multiple changes in climatic impact-drivers. Changes in several climatic impact-drivers would be more widespread at 2°C compared to 1.5°C global warming and even more widespread and/or pronounced for higher warming levels.
- C.3 Low-likelihood outcomes, such as ice sheet collapse, abrupt ocean circulation changes, some compound extreme events and warming substantially larger than the assessed very likely range of future warming cannot be ruled out and are part of risk assessment.


# D. Limiting Future Climate Change

D.1 From a physical science perspective, limiting human-induced global warming to a specific level requires limiting cumulative CO<sub>2</sub> emissions, reaching at least net zero CO<sub>2</sub> emissions, along with strong reductions in other greenhouse gas emissions. Strong, rapid and sustained reductions in CH<sub>4</sub> emissions would also limit the warming effect resulting from declining aerosol pollution and would improve air quality.


# https://interactive-atlas.ipcc.ch








Q & + m 0 0 = 10 " » :



ovrf.umd.edu v C N v symulacje klimatu

# CWRF Climate - Weather Research and Forecasting Model

Monte | Devidence | Repairs | Foreign | Seriemence | Application | Fratision | Many | Rubberton




- CWRF is a Climate extension of the Weather Research and Forecasting model (WRF):
  - → Inherits all WRF functionalities for NWP while enhancing the capability to predict climate, thus has unified applications for both weather forecast and climate prediction.
- CWRF incorporates a grand ensemble of alternative physics schemes:
  - → Contains more than 10<sup>24</sup> of alternative physics configurations representing interactions between surface

#### News:

- → CWRF test data available for download
- CWRF website goes live
- CWRF V3.1.1 ready to release

#### Links:

- CAR: Cloud-Aerosol-Radiation Ensemble Modeling System
- WRF: The Weather Research and Forecasting Model
- ☑ ESSIC/UMD



Purchase

Download



Community

Software



Support

Contact Us

## Software update

Software update

#### ANNOUNCEMENTS:

Home

EdGCM 4.0 beta is now available for Mac OS X Lion (10.7) and Mountain Lion (10.8) read more...

EdGCM 3.2 users: Windows 7 and Mac OS X 10.6 (Snow Leopard) updates are still available read more...

EdGCM provides a research-grade Global Climate Model (GCM) with a user-friendly interface that can be run on a desktop computer. For the first time, students can explore the subject of climate change in the same way that actual research scientists do. In the process of using EdGCM, students will become knowledgeable about a topic that will surely affect their lives, and we will better prepare the next generation of scientists who will grapple with a myriad of complex climate issues.

Our goal is to improve the quality of teaching and learning of climate-change science through broader access to GCMs, and to provide appropriate technology and materials to help educators use these models effectively. With research-quality resources in place, linking classrooms to actual research projects is not only possible, but can also be beneficial to the education and research communities alike.



Read more...

