

Global warming - physicist's perspective – 03

Natural history of climate

Szymon P. Malinowski


University of Warsaw, Faculty of Physics, Institute of





NAUKA O KLIMACIE
DLA SCEPTYCZNYCH



UNIVERSITY
OF WARSAW



Changes in atmospheric composition



Paleoclimatic data

Dating methods

1.1 Radiometric dating

1.2 Fission-track dating

1.3 Cosmogenic nuclide

geochronology

1.4 Luminescence dating

1.5 Incremental dating


1.6 Paleomagnetic dating

1.7 Magnetostratigraphy

1.8 Chemostratigraphy

1.9 Correlation of marker

horizons



Radiometric dating

TABLE 3-1 Radioactive Decay Used to Date Climate Records

Parent isotope	Daughter isotope	Half-life	Useful for ages:	Useful for dating:
Rubidium-87 (⁸⁷ Rb)	Strontium-87 (⁸⁷ Sr)	47 Byr	100 Myr	Granites
Uranium-238 (²³⁸ U)	Lead-206 (²⁰⁶ Pb)	4.5 Byr	>100 Myr	Many rocks
Uranium-235 (²³⁵ U)	Lead-207 (²⁰⁷ Pb)	0.7 Byr	>100 Myr	Many rocks
Potassium-40 (⁴⁰ K)	Argon-40 (⁴⁰ Ar)	1.3 Byr	>100,000 years	Basalts
Thorium 230 (²³⁰ Th)	Radon-226* (²²⁶ Ra)	75,000 years	<400,000 years	Corals
Carbon-14 (¹⁴ C)	Nitrogen-14* (¹⁴ N)	5,780 years	<50,000 years	Anything that contains carbon


**Daughter is a gas that has escaped and cannot be measured.*

Isotope systematics in the hydrological cycle:

The isotope composition of natural meteoric waters (which form the main reservoirs of freshwater on earth, ice caps, lakes, rivers and groundwater) is determined by three main factors.

- 1) The isotopic composition of the source of the moisture i.e. the ocean, the largest water reservoir on earth.
- 2) Processes of fractionation during evaporation from the ocean.
- 3) Fractionation processes during condensation in the cloud and precipitation to the ground.

$\delta^{18}\text{O}$ - Tracer of the water cycle




$\delta^{18}\text{O}$ signal in different climate archives


- H_2^{16}O

Abundance:


- $^{16}\text{O} = 99.7621$
 $^{17}\text{O} = 0.03790$
 $^{18}\text{O} = 0.20004$
- $\text{H} = 99.9844$
 $\text{D} = 0.01557$




Reduced amount of carbon stable isotope ^{13}C in plants depending on photosynthesis type.



Long-term $\delta^{18}\text{O}$ evolution



Zachos et al,
2001, Science,
v.292, p.686



Westerhold, T., N Marwan, AJ Drury, D Liebrand, C Agnini, E Anagnostou, JSK Barnet, SM Bohaty, D Vleeschouwer, F Florindo, T Frederichs, DA Hodell, AE Holbourn, D Kroon, V Lauretano, K Littler, LJ Lourens, M Lyle, H Pälike, U Röhl, J Tian, RH Wilkens, PA Wilson, JC Zachos, 2020, An astronomically dated record of Earth's climate and its predictability over the last 66 Million Years, *Science*, v. 369, Issue 6509, pp. 1383-1387. DOI: 10.1126/science.aba6853






Fig. 2. Climate states of the Cenozoic. Deep-sea benthic foraminifer high-resolution carbon (A) and oxygen (B) isotope records and the respective recurrence plots as well as scatterplots of long-term benthic foraminifer carbon versus oxygen values (C) and oxygen values versus atmospheric CO_2 concentrations (D).








Fig. 4. The relationship between Phanerozoic temperature and atmospheric CO₂.


(A) PhanDA GMST (top) and reconstructed atmospheric CO₂ (bottom), resolved at the stage level. The CO₂ reconstruction (29) is largely based on the data from Foster et al. (2017) (78) in the Paleozoic and Mesozoic, and the data from Rae et al. (2021) (80) in the Cenozoic. Shading reflects percentiles.

(B) PhanDA GMST versus CO₂, color-coded by geologic era. The York regression (86), which accounts for uncertainty in both the predictor and response variables, is shown by the black dashed line.


(C) CO₂ ranges for each of the climate states defined in Fig. 3. Light and dark bands show the 5th to 95th and 16th to 84th percentiles, respectively. The thick solid line shows the median value, and the dashed line shows the median, excluding data from the Mesozoic, where CO₂ is more uncertain.

- Long-term (> 10s of millions of years to 4.6Gyr) climate history information can come from geology.
- Oxygen isotopes ($\delta^{18}\text{O}$) from carbonate (CaCO_3) can be used as a paleotemperature proxy.
- Can also use similar techniques to measure ancient carbon dioxide levels.
- Earth has only had polar ice for ~15% of its history; frequently there is sufficient equator-to-pole heat transport to allow palm trees at the poles.
- Specific example: Snowball Earth, when Earth froze over 600 Myr ago.
- Snowball Earths are reversed by build up of carbon dioxide, and are followed by global hothouses when the ice melts.
- Earth's climate history is a total roller coaster on 10 million year to billion year timescales.

Abrupt Climate Change – example: PETM



The Palaeocene-Eocene Thermal Maximum as recorded in benthic (bottom dwelling) foraminifer (*Nuttallides truempyi*) isotopic records from sites in the Antarctic, south Atlantic and Pacific (see Zachos et al., 2003 for details). The rapid decrease in carbon isotope ratios in the top panel is indicative of a large increase in atmospheric greenhouse gases CO_2 and CH_4 that was coincident with an approximately 5°C global warming (centre panel). Using the carbon isotope records, numerical models show that CH_4 released by the rapid decomposition of marine hydrates might have been a major component ($\sim 2,000 \text{ GtC}$) of the carbon flux (Dickens and Owen, 1996).



MIS - marine Isotopic Stages, minima of $\delta^{18}\text{O}$



nature > nature communications > articles > article

Article | [Open access](#) | Published: 16 August 2019

The configuration of Northern Hemisphere ice sheets through the Quaternary

Christine L. Batchelor , Martin Maroosi, Mario Knapp, Della K. Murton, April S. Dalton, Philip L. Gibbard, Chris R. Stokes, Julian B. Murton & Andrea Manica

Nature Communications 10, Article number: 3713 (2019) | [Cite this article](#)




Comparison of NH ice-sheet extent during the last glacial cycle and MIS 6.

a shows a comparison of the reconstructed ice-sheet extent during the LGM and MIS 4.

The orange fill shows areas that were covered by ice sheets during both the LGM and MIS 4.


b shows a comparison of the reconstructed geographical maximum ice-sheet extent during the last glacial cycle (MIS 2–5d) and MIS 6. The purple fill shows areas that were covered by ice sheets during both the last glacial cycle (LGC) and MIS 6. Background is ETOPO1 1 arc-minute global relief model of the Earth's surface⁷²

Post-Glacial Sea Level Rise



Carbon in climate system

(IPCC 2021)



Volcanic C Global Ca

Sustainable and Rene
Occasional Publicati

5 Present day volcanic carbon flux estimates

Terrestrial volcanism occurs within both the hydrosphere and atmosphere. Submarine and subaerial volcanism originate almost entirely within different tectonic environments (as outlined above), tapping different, although not entirely exclusive volatile sources, shown schematically in Figure 2, below.






Figure 2. Diagrammatic representation of the different volcanic environments with estimates of CO₂ emission rates and their relationship to plate tectonic environment.



Antropogenic
vs. volcanic
CO₂
emissions.

Gerlach, T. (2011): "Volcanic
Versus Anthropogenic Carbon
Dioxide", EOS, Trans. AGU,
92:24, 201-208

Fast and slow processes in the carbon cycle




Fossil fuels are not naturally a part of the fast cycle: every ton emitted changes the carbon cycle for thousands of years

Box 6.1, Table 1 | The main natural processes that remove CO₂ consecutive to a large emission pulse to the atmosphere, their atmospheric CO₂ adjustment time scales, and main (bio)chemical reactions involved.



Processes	Time scale (years)	Reactions
Land uptake: Photosynthesis–respiration	1–10 ²	$6\text{CO}_2 + 6\text{H}_2\text{O} + \text{photons} \rightarrow \text{C}_6\text{H}_{12}\text{O}_6 + 6\text{O}_2$ $\text{C}_6\text{H}_{12}\text{O}_6 + 6\text{O}_2 \rightarrow 6\text{CO}_2 + 6\text{H}_2\text{O} + \text{heat}$
Ocean invasion: Seawater buffer	10–10 ³	$\text{CO}_2 + \text{CO}_3^{2-} + \text{H}_2\text{O} \rightleftharpoons 2\text{HCO}_3^-$
Reaction with calcium carbonate	10 ³ –10 ⁴	$\text{CO}_2 + \text{CaCO}_3 + \text{H}_2\text{O} \rightarrow \text{Ca}^{2+} + 2\text{HCO}_3^-$
Silicate weathering	10 ⁴ –10 ⁶	$\text{CO}_2 + \text{CaSiO}_3 \rightarrow \text{CaCO}_3 + \text{SiO}_2$


The last two + volcanic emissions + coal/oil/gas formation from organic carbon
= slow carbon cycle

Timescales of carbon removal.




FAQ 6.2, Figure 2 | Decay of a CO₂ excess amount of 5000 PgC emitted at time zero into the atmosphere, and its subsequent redistribution into land and ocean as a function of time, computed by coupled carbon-cycle climate models. The sizes of the colour bands indicate the carbon uptake by the respective reservoir. The first two panels show the multi-model mean from a model intercomparison project (Joos et al., 2013). The last panel shows the longer term redistribution including ocean dissolution of carbonaceous sediments as computed with an Earth System Model of Intermediate Complexity (after Archer et al., 2009b).






CH₄ in climate system



N₂O in climate system



Geological Society of London Scientific Statement: what the geological record tells us about our present and future climate



Caroline H. Lear^{1*}, Pallavi Anand², Tom Blenkinsop¹, Gavin L. Foster³, Mary Gagen⁴, Babette Hoogakker⁵, Robert D. Larter⁶, Daniel J. Lunt⁷, I. Nicholas McCave⁸, Erin McClymont⁹, Richard D. Pancost¹⁰, Rosalind E.M. Rickaby¹¹, David M. Schultz¹², Colin Summerhayes¹³, Charles J.R. Williams⁷ and Jan Zalasiewicz¹⁴



Geology is the science of how the Earth functions and has evolved and, as such, it can contribute to our understanding of the climate system and how it responds to the addition of carbon dioxide (CO₂) to the atmosphere and oceans. Observations from the geological record show that atmospheric CO₂ concentrations are now at their highest levels in at least the past 3 million years. Furthermore, the current speed of human-induced CO₂ change and warming is nearly without precedent in the entire geological record, with the only known exception being the instantaneous, meteorite-induced event that caused the extinction of non-bird-like dinosaurs 66 million years ago. In short, whilst atmospheric CO₂ concentrations have varied dramatically during the geological past due to natural processes, and have often been higher than today, the current rate of CO₂ (and therefore temperature) change is unprecedented in almost the entire geological past.

The geological record shows that changes in temperature and greenhouse gas concentrations have direct impacts on sea-level, the hydrological cycle, marine and terrestrial ecosystems, and the acidification and oxygen depletion of the oceans...

The geological record provides powerful evidence that atmospheric CO₂ concentrations drive climate change, and supports multiple lines of evidence that greenhouse gases emitted by human activities are altering the Earth's climate. Moreover, the amount of anthropogenic greenhouse gases already in the atmosphere means that Earth is committed to a certain degree of warming. As the Earth's climate changes due to the burning of fossil fuels and changes in land-use, the planet we live on will experience further changes that will have increasingly drastic effects on human societies. An assessment of past climate changes helps to inform policy decisions regarding future climate change. Earth scientists will also have an important role to play in the delivery of any policies aimed at limiting future climate change.

Summary:

Modern paleoclimatology is based on physical/chemical/biological analysis of past climate footprints present in sediments and remnants from the past.

Paleoclimatology gives insight into past climates.

Paleoclimatological analysis increases our understanding of climate forcings and feedbacks, affecting distortions of carbon cycle and other biogeochemical cycles.

Alterations of carbon cycle and changes within this cycle due to feedbacks in climate system are explaining climate variability in last ~500 millions of years.